arXiv:1411.2383v2 [math.GT] 2 Nov 2016

EXPLICIT FORMULAE FOR CHERN-SIMONS INVARIANTS OF THE TWIST KNOT ORBIFOLDS AND EDGE POLYNOMIALS OF TWIST KNOTS

JI-YOUNG HAM, JOONGUL LEE

Abstract. We calculate the Chern-Simons invariants of the twist knot orbifolds using the Schläfli formula for the generalized Chern-Simons function on the family of the twist knot cone-manifold structures. Following the general instruction of Hilden, Lozano, and Montesinos-Amilibia, we here present the concrete formulae and calculations. We use the Pythagorean Theorem, which was used by Ham, Mednykh, and Petrov, to relate the complex length of the longitude and the complex distance between the two axes fixed by two generators. As an application, we calculate the Chern-Simons invariants of cyclic coverings of the hyperbolic twist knot orbifolds. We also derive some interesting results. The explicit formula of A-polynomials of twist knots are obtained from the complex distance polynomials. Hence the edge polynomials corresponding to the edges of the Newton polygons of A-polynomials of twist knots can be obtained. In particular, the number of boundary components of every incompressible surface corresponding to slope $-4n+2$ appears to be 2.

1. Introduction

In the 1970s, Chern and Simons [3] defined an invariant of a compact 3 (mod 4) - dimensional Riemannian manifold, M, which is now called the Chern-Simons invariant, $\mathrm{cs}(M)$. In the 1980s, Meyerhoff [28] extended the definition of $\mathrm{cs}(M)$ to cusped manifolds. It is the integral of a certain 3-form and an invariant of the Riemannian connection on a principal tangent bundle of M.

Various methods of finding the Chern-Simons invariant using ideal triangulations have been introduced [30, 31, 39, 6, 5, 4] and implemented [12, 15]. But, for orbifolds, to our knowledge, there does not exist a single convenient program which computes Chern-Simons invariant. The Chern-Simons invariant can also be obtained by eta-invariant, $\eta(M)$. $\mathrm{cs}(M) = \frac{3}{2}\eta(M) \pmod{\frac{1}{2}}$ [11, 38]. But it is easier to compute Chern-Simons invariants than η-invariants.

Instead of working on complicated combinatorics of 3-dimensional ideal tetrahedra to find the Chern-Simons invariants of the twist knot orbifolds, we deal with simple one dimensional singular loci. To make the computation simpler, we express the complex length of the singular locus in terms of the complex distance between the two axes fixed by two generators. To find out the complex length of the singular locus, we start from working on $\mathrm{SL}(2,C)$. The singular locus appears in $\mathrm{SL}(2,C)$ as a series of matrix multiplications. If we first calculate the complex distance and recover the complex length of the singular locus from the complex distance using

2010 *Mathematics Subject Classification.* 57M25, 51M10, 57M27, 57M50.

Key words and phrases. Chern-Simons invariant, twist knot, orbifold, A-polynomial, edge polynomial.

Lemma 3.3, the multiplication needed in $\mathrm{SL}(2,C)$ can be cut down approximately by half. Similar methods for volumes can be found in [16]. We use the Schläfli formula for the generalized Chern-Simons function on the family of a twist knot cone-manifold structures [19]. In [20] a method of calculating the Chern-Simons invariants of two-bridge knot orbifolds were introduced but without explicit formulae. Similar approaches for $SU(2)$-connections can be found in [23] and for $\mathrm{SL}(2,C)$-connections in [22]. Explicit integral formulae for Chern-Simons invariants of the Whitehead link (the two component twist link) orbifolds and their cyclic coverings are presented in [2, 1]. A brief explanation for twist knot cone-manifolds are in [16]. You can also refer to [10, 37, 24, 32, 18, 33].

The main purpose of the paper is to find the explicit and efficient formulae for Chern-Simons invariants of the twist knot orbifolds. For two-bridge hyperbolic link, there exists an angle $\alpha_0 \in [\frac{2\pi}{3}, \pi)$ for each link K such that the cone-manifold $K(\alpha)$ is hyperbolic for $\alpha \in (0, \alpha_0)$, Euclidean for $\alpha = \alpha_0$, and spherical for $\alpha \in (\alpha_0, \pi]$ [32, 18, 24, 33]. We will use the Chern-Simons invariant of the lens space $L(4n+1, 2n+1)$ calculated in [20]. The following theorem gives the formulae for T_m for even integers m. For odd integers m, we can replace T_m by T_{-m-1} as explained in Section 2. So, the following theorem actually covers all possible hyperbolic twist knots. We exclude the non-hyperbolic case, $n = 0,\ -1$.

Theorem 1.1. *Let T_{2n} be a hyperbolic twist knot. Let $T_{2n}(\alpha)$, $0 \le \alpha < \alpha_0$ be a hyperbolic cone-manifold with underlying space S^3 and with singular set T_{2n} of cone-angle α. Let k be a positive integer such that k-fold cyclic covering of $T_{2n}(\frac{2\pi}{k})$ is hyperbolic. Then the Chern-Simons invariant of $T_{2n}(\frac{2\pi}{k})$ (mod $\frac{1}{k}$ if k is even or mod $\frac{1}{2k}$ if k is odd) is given by the following formula:*

$$\begin{aligned}\mathrm{cs}\left(T_{2n}\left(\frac{2\pi}{k}\right)\right) \equiv{}& \frac{1}{2}\mathrm{cs}\left(L(4n+1,2n+1)\right)\\ &+ \frac{1}{4\pi^2}\int_{\frac{2\pi}{k}}^{\alpha_0} Im\left(2 * \log\left(M^{-2}\frac{A+iV}{A-iV}\right)\right)\, d\alpha\\ &+ \frac{1}{4\pi^2}\int_{\alpha_0}^{\pi} Im\left(\log\left(M^{-2}\frac{A+iV_1}{A-iV_1}\right) + \log\left(M^{-2}\frac{A+iV_2}{A-iV_2}\right)\right)\, d\alpha,\end{aligned}$$

where for $A = \cot\frac{\alpha}{2}$, V $(Im(V) \le 0)$, V_1, and V_2 are zeroes of the complex distance polynomial $P_{2n} = P_{2n}(V,B)$ which is either given recursively by

$$P_{2n} = \begin{cases} \left(\left(4B^4 - 8B^2 + 4\right)V^2 - 4B^4 + 8B^2 - 2\right)P_{2(n-1)} - P_{2(n-2)}, & \text{if } n > 1,\\ \left(\left(4B^4 - 8B^2 + 4\right)V^2 - 4B^4 + 8B^2 - 2\right)P_{2(n+1)} - P_{2(n+2)}, & \text{if } n < -1,\end{cases}$$

with initial conditions

$$\begin{aligned} P_{-2}(V,B) &= \left(2B^2 - 2\right)V + 2B^2 - 1,\\ P_0(V,B) &= 1,\\ P_2(V,B) &= \left(4B^4 - 8B^2 + 4\right)V^2 + \left(2 - 2B^2\right)V - 4B^4 + 6B^2 - 1\end{aligned}$$

or given explicitly by

$$P_{2n} = \begin{cases} \sum_{i=0}^{2n} \binom{n+\lfloor \frac{i}{2} \rfloor}{i} \left(2(B^2-1)(1-V)\right)^i \left(1+2(V-1)^{-1}\right)^{\lfloor \frac{1+i}{2} \rfloor} & \text{if } n \geq 0 \\ \sum_{i=0}^{-2n-1} \binom{-n+\lfloor \frac{(i-1)}{2} \rfloor}{i} \left(2(B^2-1)(V-1)\right)^i \left(1+2(V-1)^{-1}\right)^{\lfloor \frac{1+i}{2} \rfloor} & \text{if } n < 0 \end{cases}$$

where $B = \cos\frac{\alpha}{2}$ *and* V_1 *and* V_2 *approach common* V *as* α *decreases to* α_0 *and they come from the components of* V *and* $\overline{V}$*.*

We here present some derived results. Theorem 1.2 gives the recursive formulae of A-polynomial of twist knots. In [21, Theorem 1], Hoste and Shanahan presented the recursive formulae of A-polynomials of the twist knots with the opposite orientation. Theorem 1.3 gives the explicit formulae of A-polynomials of twist knots. In [26, Theorem1.1], Mathews presented the explicit formulae of A-polynomials of the twist knots with the opposite orientation. In case $n \leq 0$ of [26], there is a typo; $\left(\frac{M^2-1}{L+M^2}\right)^i$ has to be changed into $\left(\frac{1-M^2}{L+M^2}\right)^i$ [25]. For each side of slop a/b of the Newton polygon of A_{2n}, we can obtain the edge polynomial. By substituting t for L^bM^a of each term appearing along the side edge, we have the edge polynomial, $f_{a/b}(t)$. Theorem 1.4 gives the edge polynomials of twist knots. In [8, Corollary 11.5], Cooper and Long showed that the edge polynomials of a two-bridge knot are all $\pm(t-1)^k(t+1)^l$. We pin them down in case of twist knots. Corollary 1.5 tells the number of boundary components in case of slope $-4n+2$ of twist knots. From [17], we know that the number of boundary components of two-bridge knots are one or two. We pin them down in case of slope $-4n+2$ of twist knots. Proofs of derived results are in Section 3.

Theorem 1.2. *A-polynomial* $A_{2n} = A_{2n}(L,M)$ *is given recursively by*

$$A_{2n} = \begin{cases} A_u A_{2(n-1)} - M^4\left(1+LM^2\right)^4 A_{2(n-2)} & \text{if } n > 1 \\ A_u A_{2(n+1)} - M^4\left(1+LM^2\right)^4 A_{2(n+2)} & \text{if } n < -2 \end{cases}$$

with initial conditions

$$\begin{aligned} A_{-4}(L,M) &= 1 - L + 2LM^2 + 2LM^4 + L^2M^4 - L^2M^6 - LM^8 + LM^{10} + 2L^2M^{10} \\ &\quad + 2L^2M^{12} - L^2M^{14} + L^3M^{14}, \\ A_{-2}(L,M) &= 1 + LM^6, \\ A_0(L,M) &= -1, \\ A_2(L,M) &= L - LM^2 - M^4 - 2LM^4 - L^2M^4 - LM^6 + LM^8, \end{aligned}$$

where

$$A_u = 1 - L + 2LM^2 + M^4 + 2LM^4 + L^2M^4 + 2LM^6 - LM^8 + L^2M^8.$$

Theorem 1.3. *A-polynomial* $A_{2n} = A_{2n}(L,M)$ *is given explicitly by*



FIGURE 1. twist knot (left) and its mirror image (right)



FIGURE 2. Knot 6_1 with slope 2/9 (left) and with slope 5/9 (right).

$$A_{2n} = \begin{cases} -M^{2n}\left(1+LM^2\right)^{2n}\sum_{i=0}^{2n}\binom{n+\lfloor\frac{i}{2}\rfloor}{i}\left(1-M^2\right)^i\left(1+LM^2\right)^{-i} \\ \times(L-1)^{\lfloor\frac{i}{2}\rfloor}\left(LM^2-M^{-2}\right)^{\lfloor\frac{1+i}{2}\rfloor} \qquad \text{if } n\geq 0 \\ M^{-2n}\left(1+LM^2\right)^{-2n-1}\sum_{i=0}^{-2n-1}\binom{-n+\lfloor\frac{(i-1)}{2}\rfloor}{i}\left(M^2-1\right)^i\left(1+LM^2\right)^{-i} \\ \times(L-1)^{\lfloor\frac{i}{2}\rfloor}\left(LM^2-M^{-2}\right)^{\lfloor\frac{1+i}{2}\rfloor} \qquad \text{if } n<0 \end{cases}$$

Theorem 1.4. *When $n>0$, the edge polynomials of twist knots are*

$$\begin{cases} \pm(t-1) & \text{if the slope is } -4n \\ -(t-1)^n & \text{if the slope is 4 and } n \text{ is even} \\ \pm(t-1)^n & \text{if the slope is 4 and } n \text{ is odd.} \end{cases}$$

When $n<0$, the edge polynomials of twist knots are

$$\begin{cases} t+1 & \text{if the slope is } -4n+2 \\ \pm(t-1)^{-n-1} & \text{if the slope is 4 and } n \text{ is even} \\ (t-1)^{-n-1} & \text{if the slope is 4 and } n \text{ is odd.} \end{cases}$$

Corollary 1.5. *The number of boundary components of every incompressible surface corresponding to slope $-4n+2$ of twist knots are 2.*

2. Twist knots

A knot K is the twist knot if K has a regular two-dimensional projection of the form in Figure 1. K has 2 left-handed vertical crossings and m right-handed horizontal crossings. We will denote it by T_m. One can easily check that the slope

of T_m is $2/(2m+1)$ which is equivalent to the knot with slope $(m+1)/(2m+1)$ [36]. For example, Figure 2 shows two different regular projections of knot 6_1; one with slope $2/9$ (left) and the other with slope $5/9$ (right). Note that T_m and its mirror image have the same fundamental group up to orientation and hence have the same fundamental domain up to isometry in $\mathbb{H}^3$. It follows that $T_m(\alpha)$ and its mirror image have the same fundamental set upto isometry in $\mathbb{H}^3$ and have the same Chern-Simons invariant up to sign. Since the mirror image of T_m is equivalent to T_{-m-1}, when m is odd we will use T_{-m-1} for T_m. Hence a twist knot can be represented by T_{2n} for some integer n with slope $2/(4n+1)$ or $(2n+1)/(4n+1)$.

Let us denote by X_{2n} the exterior of T_{2n}. In [21], the fundamental group of X_{2n} is calculated with 2 right-handed vertical crossings as positive crossings instead of two left-handed vertical crossings. The following theorem is tailored to our purpose. The following theorem can also be obtained by reading off the fundamental group from the Schubert normal form of T_{2n} with slope $(2n+1)/(4n+1)$ [34].

Proposition 2.1.

$$\pi_1(X_{2n}) = \left\langle s, t \mid swt^{-1}w^{-1} = 1 \right\rangle,$$

where $w = (ts^{-1}t^{-1}s)^n$.

3. The complex distance polynomial and A-polynomial

Given a twist knot T_{2n} and a set of generators, $\{s, t\}$, we identifiy the set of representations, R, of $\pi_1(X_{2n})$ into $SL(2,\mathbb{C})$ with $R(\pi_1(X_{2n})) = \{(\eta(s), \eta(t)) \mid \eta \in R\}$. Since the defining relation of $\pi_1(X_{2n})$ gives the defining equation of $R(\pi_1(X_{2n}))$ [35], $R(\pi_1(X_{2n}))$ can be thought of an affine algebraic set in $\mathbb{C}^2$. $R(\pi_1(X_{2n}))$ is well-defined upto isomorphisms which arise from changing the set of generators. We say elements in R which differ by conjugations in $SL(2,\mathbb{C})$ are equivalent.

We use two coordinates to give the structure of the affine algebraic set to $R(\pi_1(X_{2n}))$. Equivalently, for some $O \in SL(2,\mathbb{C})$, we consider both η and $\eta' = O^{-1}\eta O$:

For the complex distance polynomial, we use for the coordinates

$$\eta(s) = \begin{bmatrix} (M+1/M)/2 & e^{\frac{\rho}{2}}(M-1/M)/2 \\ e^{-\frac{\rho}{2}}(M-1/M)/2 & (M+1/M)/2 \end{bmatrix},$$

$$\eta(t) = \begin{bmatrix} (M+1/M)/2 & e^{-\frac{\rho}{2}}(M-1/M)/2 \\ e^{\frac{\rho}{2}}(M-1/M)/2 & (M+1/M)/2 \end{bmatrix},$$

and for the A-polynomial,

$$\eta'(s) = \begin{bmatrix} M & 1 \\ 0 & M^{-1} \end{bmatrix}, \quad \eta'(t) = \begin{bmatrix} M & 0 \\ t & M^{-1} \end{bmatrix}.$$

In [16], the complex distance polynomial of T_{2n} is presented recursively. Theorem 3.2 gives it explicitly. It is the defining polynomial of algebraic set $R(\pi_1(X_{2n}))$ with the set of generators given in Proposition 2.1 and with the coordinates $\eta(s)$ and $\eta(t)$ in $SL(2,\mathbb{C})$. The actual computation in [16] is done after setting $M = e^{\frac{i\alpha}{2}}$ and then the variables are changed to $B = \cos\frac{\alpha}{2}$ and $V = \cosh\rho$.

Theorem 3.1. [16, Theorem 3.1] *For $B = \cos\frac{\alpha}{2}$, $V = \cosh\rho$ is a root of the following complex distance polynomial $P_{2n} = P_{2n}(V, B)$ which is given recursively by*

$$P_{2n} = \begin{cases} \left(\left(4B^4 - 8B^2 + 4\right)V^2 - 4B^4 + 8B^2 - 2\right)P_{2(n-1)} - P_{2(n-2)} & \text{if } n > 1 \\ \left(\left(4B^4 - 8B^2 + 4\right)V^2 - 4B^4 + 8B^2 - 2\right)P_{2(n+1)} - P_{2(n+2)} & \text{if } n < -1 \end{cases}$$

with initial conditions

$$\begin{aligned} P_{-2}(V, B) &= \left(2B^2 - 2\right)V + 2B^2 - 1, \\ P_0(V, B) &= 1, \\ P_2(V, B) &= \left(4B^4 - 8B^2 + 4\right)V^2 + \left(2 - 2B^2\right)V - 4B^4 + 6B^2 - 1. \end{aligned}$$

Theorem 3.2. *For $B = \cos\frac{\alpha}{2}$, $V = \cosh\rho$ is a root of the following complex distance polynomial $P_{2n} = P_{2n}(V, B)$ which is given explicitly by*

$$P_{2n} = \begin{cases} \sum_{i=0}^{2n} \binom{n+\lfloor \frac{i}{2} \rfloor}{i} \left(2(B^2-1)(1-V)\right)^i \left(1 + 2(V-1)^{-1}\right)^{\lfloor \frac{1+i}{2} \rfloor} & \text{if } n \geq 0 \\ \sum_{i=0}^{-2n-1} \binom{-n+\lfloor \frac{(i-1)}{2} \rfloor}{i} \left(2(B^2-1)(V-1)\right)^i \left(1 + 2(V-1)^{-1}\right)^{\lfloor \frac{1+i}{2} \rfloor} & \text{if } n < 0. \end{cases}$$

Proof. We write f_{2n} for the claimed formula and show that $f_{2n} = P_{2n}$. f_{2n} can be rewritten as

$$f_{2n} = \begin{cases} \sum_{j=0}^{n} \binom{n+j}{2j} \left(2(B^2-1)(V-1)\right)^{2j} \left(1 + 2(V-1)^{-1}\right)^j \\ -\sum_{j=0}^{n} \binom{n+j}{2j+1} \left(2(B^2-1)(V-1)\right)^{2j+1} \left(1 + 2(V-1)^{-1}\right)^{j+1} & \text{if } n \geq 0 \\ \sum_{j=0}^{-n-1} \binom{-n-1+j}{2j} \left(2(B^2-1)(V-1)\right)^{2j} \left(1 + 2(V-1)^{-1}\right)^j \\ +\sum_{j=0}^{-n-1} \binom{-n+j}{2j+1} \left(2(B^2-1)(V-1)\right)^{2j+1} \left(1 + 2(V-1)^{-1}\right)^{j+1} & \text{if } n < 0. \end{cases}$$

Now, the theorem follows by solving the recurrence formula with the initial conditions given in Theorem 3.1:

$$P_{2n} = \begin{cases} \left[\left(\left(2(B^2-1)(V-1)\right)^2 - 2(B^2-1)(V-1)\right)\left(1 + 2(V-1)^{-1}\right) + 1\right] \\ \times h_{n-1} - h_{n-2} \qquad \text{if } n > 1 \\ \left(2\left(B^2-1\right)(V-1)\left(1 + 2(V-1)^{-1}\right) + 1\right)h_{-n-1} - h_{-n-2} \qquad \text{if } n < -1 \end{cases}$$

where

$$\begin{aligned}
h_n &= \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n+1}{2k+1} \left(2(B^2-1)^2(V-1)^2\left(1+2(V-1)^{-1}\right)+1\right)^{n-2k} \\
&\quad \times \left(\left(2(B^2-1)^2(V-1)^2\left(1+2(V-1)^{-1}\right)+1\right)^2-1\right)^k \\
&= \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n-i}{i}(-1)^i \left[\sum_{j=0}^{n-2i} \binom{n-2i}{j} 2^{n-2i-j} \left(\left(2(B^2-1)(V-1)\right)^2\left(1+2(V-1)^{-1}\right)\right)^j\right] \\
&= \sum_{j=0}^{n} \left[\sum_{i=0}^{\lfloor \frac{n-j}{2} \rfloor} \binom{n-i}{i}\binom{n-2i}{j}(-1)^i 2^{n-2i-j}\right] \left(\left(2(B^2-1)(V-1)\right)^2\left(1+2(V-1)^{-1}\right)\right)^j \\
&= \sum_{j=0}^{n} \binom{n+1+j}{2j+1} \left(\left(2(B^2-1)(V-1)\right)^2\left(1+2(V-1)^{-1}\right)\right)^j.
\end{aligned}$$

f_{2n} can be obtained by simplifying the above formula. □

proof of Theorem 1.2 and Theorem 1.3. Using Theorem 4.4 in [16], we get the following Lemma 3.3 which relates the zeroes of the complex distance polynomial, $P_{2n} = P_{2n}(V,B)$, and the zeroes of the A-polynomial, $A_{2n} = A_{2n}(L,M)$. Theorem 1.2 (resp. Theorem 1.3) can be obtained from P_{2n} of Theorem 3.1 (resp. Theorem 3.2) by replacing V with $\left((M^2+1)(LM^2-1)\right)/\left((M^2-1)(LM^2+1)\right)$ using the equality in Lemma 3.3 and B with $\left(M+M^{-1}\right)/2$ and by clearing denominators. □

Lemma 3.3.

$$(1) \qquad iV = A\frac{LM^2-1}{LM^2+1} \quad \textit{and} \quad L = M^{-2}\frac{A+iV}{A-iV},$$

$$(2) \qquad V = \frac{(M^2+1)(LM^2-1)}{(M^2-1)(LM^2+1)} \quad \textit{and} \quad L = M^{-2}\frac{VM^2-V+M^2+1}{-VM^2+V+M^2+1}.$$

Proof. (1) is Theorem 4.4 in [16].

Since

$$\frac{A}{i} = \frac{\cos\frac{\alpha}{2}}{i\sin\frac{\alpha}{2}} = \frac{\cosh\frac{i\alpha}{2}}{\sinh\frac{i\alpha}{2}} = \frac{\frac{M+M^{-1}}{2}}{\frac{M-M^{-1}}{2}} = \frac{M^2+1}{M^2-1},$$

we get the first equality of (2). By solving the first equality for L, we get the second equality of (2). □

Proof of Theorem 1.4. By Lemma 3.4, we have Newton polygons, NP, associated to A_{2n} in Figure 3. Let us only consider nonzero slopes if $n \neq -1$.

When $n > 0$, NP has two slopes $-4n$ and 4. When the slope is $-4n$ and the edge has the term $-M^{4n}$ on it, there are only two terms of A_{2n} appearing along the edge of the slope. From the explicit formula of A_{2n} of Theorem 1.3, they are $-M^{4n}$ and L. The term $-M^{4n}$ occurs when $i = 2n$. We get the term L by adding two terms $(n-1)L$ which occurs when $i = 2n-1$ and $-nL$ which occurs when $i = 2n$. From $L - M^{4n}$ by substituting Lt for M^{4n} and dividing by L, we get the

edge polynomial $1-t$. Hence, on the right above edge with the same slope $-4n$, we have $t-1$ for the edge polynomial. When the slope is 4 and the term $-M^{4n}$ is on the edge, from the explicit formula of A_{2n} of Theorem 1.3, nLM^{4n+4} is on the edge and the term occurs when $i=2n$. We can get the edge polynomial from the sum of the terms which appear on this edge by substituting t for LM^4 and dividing by M^{4n}. Hence the coefficient of LM^{4n+4} is the coefficient of t of the edge polynomial. Since we know the constant term is -1 and the coefficient of t is n, we get the edge polynomial $(t-1)^n$ when n is odd and $-(t-1)^n$ when n is even because the fact that the edge polynomials of two-bridge knots are up to sign the product of some powers of $t-1$ and some powers of $t+1$ [8] and the coefficient conditions forces the power of $t+1$ to be zero. Hence, on the right below edge with the same slope 4, we have $-(t-1)^n$ for the edge polynomial.

When $n<0$, NP has two slopes $-4n+2$ and 4. When the slope is $-4n+2$ and the edge has the term 1 on it, there are only two terms of A_{2n} appearing along the edge of the slope. From the explicit formula of A_{2n} of Theorem 1.3, they are 1 and LM^{-4n+2}. The term 1 occurs when $i=2n$. We get the term LM^{-4n+2} by adding two terms $(n+1)LM^{-4n+2}$ which occurs when $i=-2n-2$ and $-nLM^{-4n+2}$ which occurs when $i=-2n-1$. From $1+LM^{-4n+2}$ by substituting t for LM^{-4n+2}, we get the edge polynomial $t+1$. Hence, on the right above edge with the same slope $-4n+2$, we have $t+1$ for the edge polynomial. When the slope is 4 and the term LM^{-4n+2} is on the edge, from the explicit formula of A_{2n} of Theorem 1.3, $(n+1)L^2M^{-4n+6}$ is on the edge. We get the term $(n+1)L^2M^{-4n+6}$ by adding two terms $\frac{(n+1)(n+2)}{2}L^2M^{-4n+6}$ which occurs when $i=-2n-2$ and $\frac{-n(n+1)}{2}L^2M^{-4n+6}$ which occurs when $i=-2n-1$. We can get the edge polynomial from the sum of the terms which appear on this edge by substituting t for LM^4 and dividing by LM^{-4n+2}. Hence the coefficient of L^2M^{-4n+6} is the coefficient of t of the edge polynomial. Since we know the constant term is 1 and the coefficient of t is $n+1$, we get the edge polynomial $(t-1)^{-n-1}$ when n is odd and $-(t-1)^{-n-1}$ when n is even because the fact that the edge polynomials of two-bridge knots are up to sign the product of some powers of $t-1$ and some powers $t+1$ [8] and the coefficient conditions forces the power of $t+1$ to be zero. Hence, on the right below edge with the same slope 4, we have $(t-1)^{-n-1}$ for the edge polynomial. □

Lemma 3.4. *The Newton polygons associated to A_{2n} are polygons in Figure 3.*

Proof. The lemma is true when $n=-2$, -1, or 1.

Since the Newton polygon of A_{2n} has ones on the corners up to sign [9], to determine the shape of the Newton polygon, we only need to consider A_{2n} modulo 2. We will use the recursive formula of A_{2n} of Theorem 1.2. In modulo 2, A_u has 6 terms and $M^4\left(1+LM^2\right)^4$ has two terms $M^4+L^4M^{12}$. The lemma can be proved by induction. You just have to combine six copies of Newton polygons of A_{2n-1} (if $n>1$) or A_{2n+1} (if $n<-2$) shifted by A_u and two copies of Newton polygons of A_{2n-2} (if $n>1$) or A_{2n+1} (if $n<-2$) shifted by $M^4+L^4M^{12}$ removing double points. □

Proof of Corollary 1.5. The number of boundary components of two-bridge knots are one or two [17]. Hence the number of boundary components of every incompressible surface corresponding to slope $-4n+2$ are bounded above by 2.



FIGURE 3. Newton polygons of A_{2n}.

The orders of roots, roots of unity, of $f_{a/b}$ divide the number of boundary components of every incompressible surface corresponding to slope a/b [7]. Hence, when the slope is $-4n+2$, since there is a single root of unity of degree 2 by Theorem 1.4, the number of boundary components of every incompressible surface corresponding to slope $-4n+2$ is bounded below by 2. □

4. Generalized Chern-Simons function

The general references for this section are [19, 20, 38] and [29]. We introduce the generalized Chern-Simons function on the family of a twist knot cone-manifold structures. For the oriented knot T_{2n}, we orient a chosen meridian s such that the orientation of s followed by orientation of T_{2n} coincides with orientation of S^3. Hence, we use the definition of Lens space in [20] so that we can have the right orientation when the definition of Lens space is combined with the following frame field. On the Riemannian manifold $S^3 - T_{2n} - s$ we choose a special frame field Γ. A *special* frame field $\Gamma = (e_1, e_2, e_3)$ is an orthonomal frame field such that for each point x near T_{2n}, $e_1(x)$ has the knot direction, $e_2(x)$ has the tangent direction of a meridian curve, and $e_3(x)$ has the knot to point direction. A special frame field always exists by the proposition 3.1 of [19]. From Γ we obtain an orthonomal frame field Γ_α on $T_{2n}(\alpha) - s$ by the Schmidt orthonormalization process with respect to the geometric structure of the cone manifold $T_{2n}(\alpha)$. Moreover it can be made special by deforming it in a neighborhood of the singular set and s if necessary. Γ' is an extention of Γ to $S^3 - T_{2n}$. For each cone-manifold $T_{2n}(\alpha)$, we assign the real number:

$$I\left(T_{2n}(\alpha)\right) = \frac{1}{2}\int_{\Gamma(S^3 - T_{2n} - s)} Q - \frac{1}{4\pi}\tau(s, \Gamma') - \frac{1}{4\pi}\left(\frac{\beta\alpha}{2\pi}\right),$$

where $-2\pi \leq \beta \leq 2\pi$, Q is the Chern-Simons form:

$$Q = \frac{1}{4\pi^2}\left(\theta_{12}\wedge\theta_{13}\wedge\theta_{23} + \theta_{12}\wedge\Omega_{12} + \theta_{13}\wedge\Omega_{13} + \theta_{23}\wedge\Omega_{23}\right),$$

and

$$\tau(s,\Gamma') = -\int_{\Gamma'(s)}\theta_{23},$$

where (θ_{ij}) is the connection 1-form, (Ω_{ij}) is the curvature 2-form of the Riemannian connection on $T_{2n}(\alpha)$ and the integral is over the orthonomalizations of the same frame field. When $\alpha = \frac{2\pi}{k}$ for some positive integer, $I\left(T_{2n}\left(\frac{2\pi}{k}\right)\right)$ (mod $\frac{1}{k}$ if k is even or mod $\frac{1}{2k}$ if k is odd) is independent of the frame field Γ and of the representative in the equivalence class $\overline{\beta}$ and hence an invariant of the orbifold $T_{2n}\left(\frac{2\pi}{k}\right)$. $I\left(T_{2n}\left(\frac{2\pi}{k}\right)\right)$ (mod $\frac{1}{k}$ if k is even or mod $\frac{1}{2k}$ if k is odd) is called *the Chern-Simons invariant of the orbifold* and is denoted by $\mathrm{cs}\left(T_{2n}\left(\frac{2\pi}{k}\right)\right)$.

On the generalized Chern-Simons function on the family of the twist knot cone-manifold structures we have the following Schläfli formula.

Theorem 4.1. *(Theorem 1.2 of* [20]*) For a family of geometric cone-manifold structures, $T_{2n}(\alpha)$, and differentiable functions $\alpha(t)$ and $\beta(t)$ of t we have*

$$dI\left(T_{2n}(\alpha)\right) = -\frac{1}{4\pi^2}\beta d\alpha.$$

5. Proof of the theorem 1.1

For $n \geq 1$ and $M = e^{i\frac{\alpha}{2}}$ $(B = \cos\frac{\alpha}{2})$, $A_{2n}(M,L)$ and $P_{2n}(V,A)$ have $2n$ component zeros, and for $n < -1$, $-(2n+1)$ component zeros. The component which gives the maximal volume is the geometric component [16, 13, 14] and in [16] it is identified. For each T_{2n}, there exists an angle $\alpha_0 \in [\frac{2\pi}{3}, \pi)$ such that T_{2n} is hyperbolic for $\alpha \in (0, \alpha_0)$, Euclidean for $\alpha = \alpha_0$, and spherical for $\alpha \in (\alpha_0, \pi]$ [32, 18, 24, 33]. Denote by $D(T_{2n}(\alpha))$ be the set of common zeros of the discriminant of $A_{2n}(L, e^{\frac{i\alpha}{2}})$ over L and the discriminant of $P_{2n}(V, \cos\frac{\alpha}{2})$ over V. Then α_0 will be one of $D(T_{2n}(\alpha))$.

On the geometric component we can calculate the Chern-Simons invariant of an orbifold $T_{2n}(\frac{2\pi}{k})$ (mod $\frac{1}{k}$ if k is even or mod $\frac{1}{2k}$ if k is odd), where k is a positive

integer such that k-fold cyclic covering of $T_{2n}(\frac{2\pi}{k})$ is hyperbolic:

$$\begin{aligned}
\mathrm{cs}\left(T_{2n}\left(\frac{2\pi}{k}\right)\right) &\equiv I\left(T_{2n}\left(\frac{2\pi}{k}\right)\right) \qquad \left(\mathrm{mod}\ \frac{1}{k}\right)\\
&\equiv I\left(T_{2n}(\pi)\right) + \frac{1}{4\pi^2}\int_{\frac{2\pi}{k}}^{\pi} \beta\, d\alpha \qquad \left(\mathrm{mod}\ \frac{1}{k}\right)\\
&\equiv \frac{1}{2}\mathrm{cs}\left(L(4n+1,2n+1)\right)\\
&+ \frac{1}{4\pi^2}\int_{\frac{2\pi}{k}}^{\alpha_0} Im\left(2 * \log\left(M^{-2}\frac{A+iV}{A-iV}\right)\right)\, d\alpha\\
&+ \frac{1}{4\pi^2}\int_{\alpha_0}^{\pi} Im\left(\log\left(M^{-2}\frac{A+iV_1}{A-iV_1}\right) + \log\left(M^{-2}\frac{A+iV_2}{A-iV_2}\right)\right)\, d\alpha\\
&\left(\mathrm{mod}\ \frac{1}{k} \text{ if } k \text{ is even or mod } \frac{1}{2k} \text{ if } k \text{ is odd}\right)
\end{aligned}$$

where the second equivalence comes from Theorem 4.1 and the third equivalence comes from the fact that $I\left(T_{2n}(\pi)\right) \equiv \frac{1}{2}\mathrm{cs}\left(L(4n+1,2n+1)\right)\left(\mathrm{mod}\ \frac{1}{2}\right)$, Lemma 3.3, and geometric interpretations of hyperbolic and spherical holonomy representations.

The fundamental set of the two-bridge link orbifolds are constructed in [27]. The following theorem gives the Chern-Simons invariant of the Lens space $L(4n+1, 2n+1)$.

Theorem 5.1. *(Theorem 1.3 of* [20]*)*

$$\mathrm{cs}\left(L\left(4n+1,2n+1\right)\right) \equiv \frac{6n+4}{8n+2} \qquad (mod\ 1).$$

6. Chern-Simons invariants of the hyperbolic twist knot orbifolds and of their cyclic coverings

The table 1 (resp. the table 2) gives the approximate Chern-Simons invariant of the hyperbolic twist knot orbifold, $\mathrm{cs}\left(T_{2n}(\frac{2\pi}{k})\right)$ for n between 2 and 9 (resp. for n between -9 and -2) and for k between 3 and 10, and of its cyclic covering, $cs\left(M_k(T_{2n})\right)$. We used Simpson's rule for the approximation with 10^4 (5×10^3 in Simpson's rule) intervals from $\frac{2\pi}{k}$ to α_0 and 10^4 (5×10^3 in Simpson's rule) intervals from α_0 to π. The table 3 gives the approximate Chern-Simons invariant of T_{2n} for each n between -9 and 9 except the unknot, the torus knot, and the amphicheiral knot. We again used Simpson's rule for the approximation with 10^4 (5×10^3 in Simpson's rule) intervals from 0 to α_0 and 10^4 (5×10^3 in Simpson's rule) intervals from α_0 to π. We used Mathematica for the calculations. We record here that our data in table 3 and those obtained from SnapPy match up up to six decimal points.

Acknowledgements. The authors would like to thank Alexander Mednykh and Hyuk Kim for their various helps, Nathan Dunfield and Daniel Mathews for their prompt helps and anonymous referees.

References

[1] N. V. Abrosimov. The Chern-Simons invariants of cone-manifolds with the Whitehead link singular set [translation of mr2485364]. *Siberian Adv. Math.*, 18(2):77–85, 2008.

[2] Nikolay V. Abrosimov. On Chern-Simons invariants of geometric 3-manifolds. *Sib. Èlektron. Mat. Izv.*, 3:67–70 (electronic), 2006.

Table 1: Chern-Simons invariant of the hyperbolic twist knot orbifold, $\mathrm{cs}\left(T_{2n}(\frac{2\pi}{k})\right)$ for n between 2 and 9 and for k between 3 and 10, and of its cyclic covering, $\mathrm{cs}\left(M_k(T_{2n})\right)$.

k	$\mathrm{cs}\left(T_4(\frac{2\pi}{k})\right)$	$\mathrm{cs}\left(M_k(T_4)\right)$
3	0.0875301	0.262590
4	0.144925	0.579699
5	0.0784576	0.392288
6	0.0351571	0.210943
7	0.00506505	0.0354553
8	0.108039	0.864313
9	0.0218112	0.196301
10	0.0530574	0.530574

k	$\mathrm{cs}\left(T_6(\frac{2\pi}{k})\right)$	$\mathrm{cs}\left(M_k(T_6)\right)$
3	0.0449535	0.134860
4	0.0876043	0.350417
5	0.0165337	0.0826684
6	0.138167	0.829004
7	0.0120078	0.0840545
8	0.0430876	0.344700
9	0.0121250	0.109125
10	0.0876213	0.876213

k	$\mathrm{cs}\left(T_8(\frac{2\pi}{k})\right)$	$\mathrm{cs}\left(M_k(T_8)\right)$
3	0.0161266	0.0483799
4	0.0536832	0.214733
5	0.0817026	0.408513
6	0.103012	0.618074
7	0.0481239	0.336867
8	0.00768503	0.0614802
9	0.0322210	0.289989
10	0.0521232	0.521232

k	$\mathrm{cs}\left(T_{10}(\frac{2\pi}{k})\right)$	$\mathrm{cs}\left(M_k(T_{10})\right)$
3	0.162697	0.488091
4	0.0320099	0.128040
5	0.0597580	0.298790
6	0.0809665	0.485799
7	0.0260276	0.182193
8	0.110559	0.884475
9	0.0100766	0.0906893
10	0.0299660	0.299660

k	$\mathrm{cs}\left(T_{12}(\frac{2\pi}{k})\right)$	$\mathrm{cs}\left(M_k(T_{12})\right)$
3	0.148360	0.445081
4	0.0170833	0.0683334
5	0.0447221	0.223610
6	0.0658884	0.395330
7	0.0109281	0.0764969
8	0.0954474	0.763579
9	0.0505121	0.454609
10	0.0148406	0.148406

k	$\mathrm{cs}\left(T_{14}(\frac{2\pi}{k})\right)$	$\mathrm{cs}\left(M_k(T_{14})\right)$
3	0.137750	0.413249
4	0.00620422	0.0248169
5	0.0337900	0.168950
6	0.0549355	0.329613
7	0.0713932	0.499752
8	0.0844779	0.675823
9	0.0395376	0.355839
10	0.00386414	0.0386414

k	$\mathrm{cs}\left(T_{16}(\frac{2\pi}{k})\right)$	$\mathrm{cs}\left(M_k(T_{16})\right)$
3	0.129617	0.388850
4	0.247931	0.991725
5	0.0254880	0.127440
6	0.0466221	0.279733
7	0.0630741	0.441518
8	0.0761526	0.609221
9	0.0312096	0.280887
10	0.0955375	0.955375

k	$\mathrm{cs}\left(T_{18}(\frac{2\pi}{k})\right)$	$\mathrm{cs}\left(M_k(T_{18})\right)$
3	0.123198	0.369593
4	0.241431	0.965725
5	0.0189709	0.0948543
6	0.0400981	0.240588
7	0.0565432	0.395803
8	0.0696194	0.556955
9	0.0246862	0.222176
10	0.0890057	0.890057

[3] Shiing-shen Chern and James Simons. Some cohomology classes in principal fiber bundles and their application to Riemannian geometry. *Proc. Nat. Acad. Sci. U.S.A.*, 68:791–794, 1971.

Table 2: Chern-Simons invariant of the hyperbolic twist knot orbifold, $\mathrm{cs}\left(T_{2n}(\frac{2\pi}{k})\right)$ for n between -9 and -2 and for k between 3 and 10, and of its cyclic covering, $\mathrm{cs}\left(M_k(T_{2n})\right)$.

k	$\mathrm{cs}\left(T_{-4}(\frac{2\pi}{k})\right)$	$\mathrm{cs}\left(M_k(T_{-4})\right)$
3	0.0200144	0.0600431
4	0.186811	0.747246
5	0.00166667	0.00833333
6	0.0504622	0.302773
7	0.0163442	0.114410
8	0.116990	0.935921
9	0.0292902	0.263612
10	0.0595432	0.595432

k	$\mathrm{cs}\left(T_{-6}(\frac{2\pi}{k})\right)$	$\mathrm{cs}\left(M_k(T_{-6})\right)$
3	0.0749433	0.224830
4	0.0126376	0.0505506
5	0.0873477	0.436738
6	0.140792	0.844753
7	0.0376998	0.263898
8	0.0862114	0.689691
9	0.0133130	0.119817
10	0.0552937	0.552937

k	$\mathrm{cs}\left(T_{-8}(\frac{2\pi}{k})\right)$	$\mathrm{cs}\left(M_k(T_{-8})\right)$
3	0.109659	0.328978
4	0.0564153	0.225661
5	0.0330919	0.165460
6	0.0205610	0.123366
7	0.0130371	0.0912595
8	0.00816423	0.0653138
9	0.00482762	0.0434486
10	0.00244289	0.0244289

k	$\mathrm{cs}\left(T_{-10}(\frac{2\pi}{k})\right)$	$\mathrm{cs}\left(M_k(T_{-10})\right)$
3	0.133696	0.401087
4	0.0832469	0.332988
5	0.0603968	0.301984
6	0.0480386	0.288232
7	0.0405999	0.284200
8	0.0357763	0.286210
9	0.0324710	0.292239
10	0.0301075	0.301075

k	$\mathrm{cs}\left(T_{-12}(\frac{2\pi}{k})\right)$	$\mathrm{cs}\left(M_k(T_{-12})\right)$
3	0.150597	0.451792
4	0.101087	0.404347
5	0.0784041	0.392020
6	0.0661095	0.396657
7	0.0587029	0.410920
8	0.0538979	0.431183
9	0.0506045	0.455441
10	0.0482492	0.482492

k	$\mathrm{cs}\left(T_{-14}(\frac{2\pi}{k})\right)$	$\mathrm{cs}\left(M_k(T_{-14})\right)$
3	0.162874	0.488621
4	0.113753	0.455011
5	0.0911448	0.455724
6	0.0788794	0.473276
7	0.0000589596	0.000412717
8	0.0666914	0.533532
9	0.00784780	0.0706302
10	0.0610519	0.610519

k	$\mathrm{cs}\left(T_{-16}(\frac{2\pi}{k})\right)$	$\mathrm{cs}\left(M_k(T_{-16})\right)$
3	0.00545933	0.0163780
4	0.123196	0.492785
5	0.000626850	0.00313425
6	0.0883767	0.530260
7	0.00956397	0.0669478
8	0.0762009	0.609607
9	0.0173633	0.156270
10	0.0705667	0.705667

k	$\mathrm{cs}\left(T_{-18}(\frac{2\pi}{k})\right)$	$\mathrm{cs}\left(M_k(T_{-18})\right)$
3	0.0126607	0.0379822
4	0.130503	0.522012
5	0.00795573	0.0397786
6	0.0957144	0.574286
7	0.0169085	0.118359
8	0.0835496	0.668397
9	0.0247059	0.222353
10	0.0779144	0.779144

[4] Jinseok Cho, Hyuk Kim, and Seonwha Kim. Optimistic limits of kashaev invariants and complex volumes of hyperbolic links. *J. Knot Theory Ramifications*, 23(9), 2014.

[5] Jinseok Cho and Jun Murakami. The complex volumes of twist knots via colored Jones polynomials. *J. Knot Theory Ramifications*, 19(11):1401–1421, 2010.

Table 3: Chern-Simons invariant of T_{2n} for n between 2 and 9 and for n between -9 and -2).

2n	α_0	cs (T_{2n})
4	2.57414	0.344023
6	2.75069	0.277867
8	2.84321	0.242222
10	2.90026	0.220016
12	2.93897	0.204869
14	2.96697	0.193882
16	2.98817	0.185550
18	3.00477	0.179014

2n	α_0	cs (T_{2n})
-4	2.40717	0.346796
-6	2.67879	0.444846
-8	2.80318	0.492293
-10	2.87475	0.0200385
-12	2.92130	0.0382117
-14	2.95401	0.0510293
-16	2.97825	0.0605519
-18	2.99694	0.0679043

[6] Jinseok Cho, Jun Murakami, and Yoshiyuki Yokota. The complex volumes of twist knots. *Proc. Amer. Math. Soc.*, 137(10):3533–3541, 2009.

[7] D. Cooper, M. Culler, H. Gillet, D. D. Long, and P. B. Shalen. Plane curves associated to character varieties of 3-manifolds. *Invent. Math.*, 118(1):47–84, 1994.

[8] D. Cooper and D. D. Long. Remarks on the A-polynomial of a knot. *J. Knot Theory Ramifications*, 5(5):609–628, 1996.

[9] D. Cooper and D. D. Long. The A-polynomial has ones in the corners. *Bull. London Math. Soc.*, 29(2):231–238, 1997.

[10] Daryl Cooper, Craig D. Hodgson, and Steven P. Kerckhoff. *Three-dimensional orbifolds and cone-manifolds*, volume 5 of *MSJ Memoirs*. Mathematical Society of Japan, Tokyo, 2000. With a postface by Sadayoshi Kojima.

[11] David Coulson, Oliver A. Goodman, Craig D. Hodgson, and Walter D. Neumann. Computing arithmetic invariants of 3-manifolds. *Experiment. Math.*, 9(1):127–152, 2000.

[12] Marc Culler, Nathan Dunfield, Jeff Weeks, and Many others. SnapPy. `http://www.math.uic.edu/t3m/SnapPy/`.

[13] Nathan M. Dunfield. Cyclic surgery, degrees of maps of character curves, and volume rigidity for hyperbolic manifolds. *Invent. Math.*, 136(3):623–657, 1999.

[14] Stefano Francaviglia and Ben Klaff. Maximal volume representations are Fuchsian. *Geom. Dedicata*, 117:111–124, 2006.

[15] Oliver Goodman. Snap. `http://sourceforge.net/projects/snap-pari`.

[16] Ji-Young Ham, Alexander Mednykh, and Vladimir Petrov. Trigonometric identities and volumes of the hyperbolic twist knot cone-manifolds. *J. Knot Theory Ramifications*, 23(12):1450064, 16, 2014.

[17] A. Hatcher and W. Thurston. Incompressible surfaces in 2-bridge knot complements. *Invent. Math.*, 79(2):225–246, 1985.

[18] Hugh Hilden, María Teresa Lozano, and José María Montesinos-Amilibia. On a remarkable polyhedron geometrizing the figure eight knot cone manifolds. *J. Math. Sci. Univ. Tokyo*, 2(3):501–561, 1995.

[19] Hugh M. Hilden, María Teresa Lozano, and José María Montesinos-Amilibia. On volumes and Chern-Simons invariants of geometric 3-manifolds. *J. Math. Sci. Univ. Tokyo*, 3(3):723–744, 1996.

[20] Hugh M. Hilden, María Teresa Lozano, and José María Montesinos-Amilibia. Volumes and Chern-Simons invariants of cyclic coverings over rational knots. In *Topology and Teichmüller spaces (Katinkulta, 1995)*, pages 31–55. World Sci. Publ., River Edge, NJ, 1996.

[21] Jim Hoste and Patrick D. Shanahan. A formula for the A-polynomial of twist knots. *J. Knot Theory Ramifications*, 13(2):193–209, 2004.

[22] Paul Kirk and Eric Klassen. Chern-Simons invariants of 3-manifolds decomposed along tori and the circle bundle over the representation space of T^2. *Comm. Math. Phys.*, 153(3):521–557, 1993.

[23] Paul A. Kirk and Eric P. Klassen. Chern-Simons invariants of 3-manifolds and representation spaces of knot groups. *Math. Ann.*, 287(2):343–367, 1990.

[24] Sadayoshi Kojima. Deformations of hyperbolic 3-cone-manifolds. *J. Differential Geom.*, 49(3):469–516, 1998.
[25] Daniel V. Mathews. Erratum: An explicit formula for the A-polynomial of twist knots [mr3268980]. *J. Knot Theory Ramifications*, 23(11):1492001, 1, 2014.
[26] Daniel V. Mathews. An explicit formula for the A-polynomial of twist knots. *J. Knot Theory Ramifications*, 23(9), 2014.
[27] Alexander Mednykh and Aleksey Rasskazov. On the structure of the canonical fundamental set for the 2-bridge link orbifolds. `www.mathematik.uni-bielefeld.de/sfb343/preprints/pr98062.ps.gz`, 1998. Universitt Bielefeld, Sonderforschungsbereich 343, Discrete Structuren in der Mathematik, Preprint, 98062.
[28] Robert Meyerhoff. Hyperbolic 3-manifolds with equal volumes but different Chern-Simons invariants. In *Low-dimensional topology and Kleinian groups (Coventry/Durham, 1984)*, volume 112 of *London Math. Soc. Lecture Note Ser.*, pages 209–215. Cambridge Univ. Press, Cambridge, 1986.
[29] Robert Meyerhoff and Daniel Ruberman. Mutation and the η-invariant. *J. Differential Geom.*, 31(1):101–130, 1990.
[30] Walter D. Neumann. Combinatorics of triangulations and the Chern-Simons invariant for hyperbolic 3-manifolds. In *Topology '90 (Columbus, OH, 1990)*, volume 1 of *Ohio State Univ. Math. Res. Inst. Publ.*, pages 243–271. de Gruyter, Berlin, 1992.
[31] Walter D. Neumann. Extended Bloch group and the Cheeger-Chern-Simons class. *Geom. Topol.*, 8:413–474 (electronic), 2004.
[32] Joan Porti. Spherical cone structures on 2-bridge knots and links. *Kobe J. Math.*, 21(1-2):61–70, 2004.
[33] Joan Porti and Hartmut Weiss. Deforming Euclidean cone 3-manifolds. *Geom. Topol.*, 11:1507–1538, 2007.
[34] Robert Riley. Parabolic representations of knot groups. I. *Proc. London Math. Soc. (3)*, 24:217–242, 1972.
[35] Robert Riley. Nonabelian representations of 2-bridge knot groups. *Quart. J. Math. Oxford Ser. (2)*, 35(138):191–208, 1984.
[36] Horst Schubert. Knoten mit zwei Brücken. *Math. Z.*, 65:133–170, 1956.
[37] William Thurston. The geometry and topology of 3-manifolds. `http://library.msri.org/books/gt3m`, 1977/78. Lecture Notes, Princeton University.
[38] Tomoyoshi Yoshida. The η-invariant of hyperbolic 3-manifolds. *Invent. Math.*, 81(3):473–514, 1985.
[39] Christian K. Zickert. The volume and Chern-Simons invariant of a representation. *Duke Math. J.*, 150(3):489–532, 2009.

Department of Science, Hongik University, 94 Wausan-ro, Mapo-gu, Seoul, 121-791, Department of Mathematical Sciences, Seoul National University, San 56-1 Shinrim-dong Kwanak-gu Seoul 151-747, Korea

E-mail address: `jiyoungham1@gmail.com.`

Department of Mathematics Education, Hongik University, 94 Wausan-ro, Mapo-gu, Seoul, 121-791, Korea

E-mail address: `jglee@hongik.ac.kr`